



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AN EASTERN RECORD AND A NOTE ON *CHARINA BOTTAE* (BLAINVILLE)

While driving through Shoshone Canyon, July 11, 1921, a *Charina* was collected four miles west of Cody, Wyoming. As far as I can determine this is the most eastern record for *Charina bottae*.

The snake was found in one of the few small open spaces along the road in the canyon. Because of the relative scarcity of specimens of this species I have given this specimen (now U. of Mich. No. 56253) to the Museum of Zoology, University of Michigan.

The Boa is like the typical *Charina bottae* except that the temporal plate is much reduced.

It is interesting to note that it differs from the specimens which Van Denburgh (Calif. Acad. Sci. Vol. X, No. 3, pp. 31-32, Aug. 6, 1920) described from Utah as *Charina bottae utahensis*. His new subspecies is based upon seven specimens each having 41 scale rows while the *C. bottae* of the more western states have from 43-49 scale rows. My specimen collected about 125 miles east of the more eastern record of *C. b. utahensis* of Van Denburgh has 43 rows. The most eastern record of the U. S. N. M. (No. 60237) from Chico, Montana, has 43 scale rows.

A. I. ORTENBURGER,
Ann Arbor, Michigan.